# THE KENYA NATIONAL EXAMINATIONS COUNCIL Kenya Certificate of Secondary Education

233/2 Paper 2

## CHEMISTRY - (Theory)

### Dec. 2022 - 2 hours



Name	Index Number
Candidate's Signature	Date

#### Instructions to candidates

- (a) Write your name and index number in the spaces provided above.
- (b) Sign and write the date of examination in the spaces provided above.
- (c) Answer all the questions in the spaces provided in the question paper.
- (d) Non-programmable silent electronic calculators and KNEC mathematical tables may be used.
- (e) All working must be clearly shown where necessary.
- (f) This paper consists of 16 printed pages.
- (g) Candidates should check the question paper to ascertain that all the pages are printed as indicated and that no questions are missing.
- (h) Candidates should answer the questions in English.

#### For Examiner's Use Only

Question	Maximum Score	Candidate's Score		
2	13			
3	12	4		
4	1.3	-Citi		
5	11	67.25		
6666		EUV		
7	I O			
Total Score	80			







1. (a) Aluminium and phosphorus form oxides with general formula M<sub>2</sub>O<sub>3</sub>. Complete **Table 1** by writing the properties of the oxides.

Table 1

Property	$Al_2O_3$	$P_2O_3$
Structure		
Bonding		
Acid/base character		

(3 marks)

(b) The grid in **Figure 1** shows part of the Periodic Table. Use it to answer the questions that follow.




Figure 1

(i) Give the total number of elements that can be placed in:

I.	period I	(½ mark)
		•••••••••••••
II.	period 5	(½ mark)

(ii) Place each of the following elements in the grid:

I.	Element X, whose atomic number is 14	(1 mark)
II.	Element Y, with the highest first ionisation energy	(1 mark)
III.	Element Z, with the lowest first ionisation energy	(1 mark)
IV.	Element L, whose ion L <sup>2-</sup> has electron arrangement 2.8	(1 mark)
V.	Element D, whose ion D2+ has electron arrangement 2.8.8	(1 mark)
VI.	Element Q, a halogen with the highest atomic radius	(1 mark)
VII.	Element R, a period 3 element that exists as a monoatomic gas	(1 mark)

2.	The general	formula	of alkanols	is $C_nH_{2n+1}$	OH.
----	-------------	---------	-------------	------------------	-----

Draw the structure and give the name of the alkanol with $n = 5$ .	(2 marks)
Structure	
***************************************	••••••
•••••••••••••••••••••••••••••••••••••••	••••••••
***************************************	••••••••
Name	

(b) Table 2 gives the boiling points of some alkanols.

Table 2

n	Boiling point/°C
2	78.5
3	97.2
4	117.0



-
0
3
2

(ii)	From the graph, determine the boiling point of the alkanol with $n = 5$ .	(1 mark)
(iii)	The boiling point of the alkanol with $n = 2$ is much higher than that of Explain ( $C = 12.0$ ; $H = 1.0$ ; $O = 16.0$ ).	butane. (2 marks)
	•••••••••••••••••••••••••••••••••••••	• • • • • • • • • • • • • • • • • • • •
		•••••••
		•••••••••••••••••••••••••••••••••••••••

	•	7	
C	•	1	

((c))	Alka	nolls ante unse	d as finel.			
	((ii))	Give anot	her use of alkamols.			(1 mark)
	((iii))	White an	equation for the comb	oustion of the alkano	l with n = 2.	(1 mark)
	((iiii))		ond energies in <b>Table</b> anol with n = 2. <b>Table</b>		nthalpy change of	combustion (3 marks)
			Bond	Energy kJ/mol		
			C - C	348		
			C – H	412		
			C - O	360		
			O - H	463		
			O = O	496		
			C = 0	743		
			FETTE			
					· · · · · · · · · · · · · · · · · · ·	••••••
			•••••		• • • • • • • • • • • • • • • • • • • •	••••••
						••••••
				***************************************		••••••
			•••••			••••••
					• • • • • • • • • • • • • • • • • • • •	•••••••
						••••••
						••••••



. (a)	Use	the following cell notation to answer the questions that follow:	
	Al(s)	$(A_{13}^{3+}(aq) // Pb^{2+}(aq) / Pb(s); E_{cell}^{\theta} = +1.53 V$	
	(i)	State what the symbol / represents.	(1 mark)
			• • • • • • • • • • • • • • • • • • • •
			• • • • • • • • • • • • • • • • • • • •
	(ii)	Write the equation for the cell reaction.	(1 mark)
			••••••
	(iii)	Given that $E^{\theta}$ value for $Pb^{2+}(aq) / Pb(s)$ is $-0.13 \text{ V}$ calculate the $E^{\theta}(aq) / Al(s)$ .	θ value for (2 marks)
		•••••••••••••••••••••••••••••••••••••••	••••••
		•••••••••••••••••••••••••••••••••••••••	••••••
			•••••••
	(iv)	State one use of electrochemical cells.	(1 mark)
			•••••••

(b) Figure 2 shows a cell used to electrolyse water.




Figure 2

(i)	State why it is necessary to add dilute sulphuric(VI) acid to the water.	(1 mark)
		•••••••
	·····	•••••••••••
		••••••••
(ii)	State the electrode at which oxygen is produced and give a reason.	(1 mark)
		••••••
	•••••••••••••••••••••••••••••••••••••••	•••••••
(iii)	Write an equation for the formation of oxygen.	(1 mark)
	•••••••••••••••••••••••••••••••••••••••	••••••

	(10)	After electrolysing the water for 88 seconds, the volume of oxygen gas conected was 23 0 cm. Determine the				
			volume of hydrogen gas collected	(I mark)		
			マニア・ア・ア・ア・ア・ア・ア・ア・ア・ア・ア・ア・ア・ア・ア・ア・ア・ア・ア・			
			ためる 我をもれたとうというという ないれるとのとれる おされる こうとうとう からかい しょうかい しょうかい しょう しゅう しゅう しゅう しゅう しゅう しゅう かんかい しょう かんしょう しょう しょう しょう しゅう しゅう しゅう しゅう しゅう しゅう しゅう しゅう しゅう しゅ			
			amount of current used (1 Faraday = 96500C)	(3 marks)		
4.	(a) Sta	te and e	splain how a catalyst affects:			
	(i)	rate	of a reaction	(2 marks)		
		******	***************************************	5.风的水面面实现的外面水面的有水面的有水面的木面的水面的		
		*****	***************************************	******************		
	(ii)	yield	d of the products	(1 mark)		
			**************************************	************		
		****	***************************************			
		*****	"世界其尊君正正是有不不可以不可以不可以不可以不可以不可以不可以不可以不可以不可以不可以不可以不可以	· · · · · · · · · · · · · · · · · · ·		

(b) Rates of reactions are measured using various methods. In the decomposition of hydrogen peroxide, the rate is measured by recording the volume of oxygen gas produced with time.

Other than measuring volume of gas produced, describe a method that can be used to measure the rates of each of the following reactions.

(i) Sodium thiosulphate with hydrochloric acid.

$S_2O_3^{2-}(aq) + 2H^+(aq) \rightarrow S(s) + SO_2(g) + H_2O(1)$	(2 marks)
•••••••••••••••••••••••••••••••••••••••	• • • • • • • • • • • • • • • • • • • •
•••••••••••••••••••••••••••••••••••••••	•••••••••••••••••••••••••••••••••••••••
•••••••••••••••••••••••••••••••••••••••	••••••••••••••••••••••••••••••••••••••

(ii) Acidified potassium manganate(VII) with iron(II) sulphate.

MnO <sub>4</sub> <sup>-</sup> (aq) +	5Fe <sup>2+</sup> (aq) +	8H+(aq) →	Mn <sup>2+</sup> (aq) +	- 5Fe <sup>3+</sup> (aq)	+ 4H <sub>2</sub> O(l) (1 mark)

In an experiment, the rate of decomposition of 50 cm<sup>3</sup> of hydrogen peroxide in the presence of manganese(IV) oxide was measured. **Figure 3** shows a graph of the results obtained.




Figure 3

(i)	Write an equation for the reaction.	(1 mark)
	••••••••••••••••••••••••••••••••••••	••••••••••••
	•••••••••••••••••••••••••••••	

(ii)	Using the graph, determine the maximum number of moles of oxygen p (Molar gas volume at room temperature and pressure = 24 dm <sup>3</sup> ).				
		(2 marks)			
	•••••••••••••••••••••••••••••••••••••••	•••••••			
	••••••••••••••••••••••••••••••••••••	••••••			
	•••••••••••••••••••••••••••••••••••	• • • • • • • • • • • • • • • • • • • •			
	•••••••••••••••••••••••••••••••••••	• • • • • • • • • • • • • • • • • • • •			
	······································				
	······································				
	***************************************				
	•••••••••••••••••••••••••••••••••••••••	• • • • • • • • • • • • • • • • • • • •			
(iii)	Calculate the concentration in moles per litre of hydrogen peroxide.	(2 marks)			
	•••••••••••••••••••••••••••••••••••••••				
	•••••••••••••••••••••••••••••••••••••••				
	•••••••••••••••••••••••••••••••••••••••				
	•••••••••••••••••••••••••••••••••••••••				
	•••••••••••••••••••••••••••••••••••••••	• • • • • • • • • • • • • • • • • • • •			
		••••••••			
	***************************************	•••••••			
<b>.</b> .		•••••••			
(iv)	Determine the rate of decomposition at the 18th second.	(1 mark)			
	•••••••••••••••••••••••••••••••••••••••	•••••••••			
	•••••••••••••••••••••••••••••••••••••••	•••••••••			
	••••••••••••••••••••••••••••••••••••••	• • • • • • • • • • • • • • • • • • • •			
	•••••••••••••••••••••••••••••••••••••••	••••••••			
	•••••••••••••••••••••••••••••••••••••••	•••••••			
(v)	State and explain one factor that would increase the rate of decompositions of the hydrogen peroxide.	tion of			
		(			
	***************************************	••••••••			
	•••••••••••••••••••••••••••••••••••••••				
	***************************************	••••••••			
	***************************************	••••••••			

5.	(a)	Expl gas a	lain how concentrated sulphuric(VI) acid can be prepared from sulphur(VI) oxi (2 n	de narks)
		•••••		
		•••••		
		•••••	······································	
	(b)	Cond	centrated sulphuric(VI) acid acts as a dehydrating and as a drying agent.	
		(i)	Give an example of a cost bot can be dried using concentrated sulphuric(VI)	acid. nark)
				••••••
		(ii)	Complete the following equations to show how concentrated sulphuric(VI) acts as a dehydrating agent.	cid
			I. $C_{12}H_{22}O_{11} \longrightarrow$ (1 n	nark)
			II. $CoCl_2 \cdot 6H_2O \longrightarrow$ (1 m	nark)
		(iii)	State the observations made when concentrated sulphuric(VI) acid dehydrates	3:
			I. $C_{12}H_{22}O_{11}$ (1 m	ark)
				•••••
				ark)
	(c)	State	the properties of concentrated sulphuric(VI) acid which are illustrated by the wing reactions:	•••••
		(i)	$S(s) + 2H_2SO_4(1) \rightarrow 3SO_2(g) + 2H_2O(1)$ (1 ma	ark)

		(ii)	$NaNO_3(s) + H_2SO_4(l) \rightarrow HNO_3(aq) + NaHSO_4(s)$	(1 mark)
	(d)	When	a mixture of 5 cm <sup>3</sup> ethanol, 1 cm <sup>3</sup> concentrated sulphuric(VI) acid and 5 coic acid was heated in a beaker, a pleasant smelling compound was forme	cm <sup>3</sup> d.
		(i)	state the role of the concentrated sulphuric(VI) acid.	(1 mark)
			•••••••••••••••••••••••••••••••••••••••	••••••
		(ii)	write the formula of the pleasant smelling compound.	(1 mark)
			•••••••••••••••••••••••••••••••••••••••	
6.		ous type ury cell	es of cells are used to electrolyse concentrated sodium chloride. One of the	m is the
	(a)	Name	e another type of cell used.	(1 mark)
	(b)		mercury cell uses titanium or graphite as anode and mercury as cathode. So is not used for the anode.	tate why (1 mark)
		•••••		•••••••
		•••••		
	(c)	At th	e anode, chloride ions and not hydroxide ions are oxidised. Give a reason.	
				•••••••
				••••••



(d)	Desci	ribe using equations, how sodium hydroxide and hydrogen are produced	in the cell. (3 marks)
	•••••	······································	•••••••••••••••••••••••••••••••••••••••
			•••••••
	•••••		••••••
	••••••	······································	•••••••
	•••••		• • • • • • • • • • • • • • • • • • • •
	•••••		•••••••
	•••••		•••••••
(e)	Give	two reasons why it is necessary to recycle the mercury used in the cell.	(2 marks)
	•••••		••••••
	•••••	•••••••••••••••••••••••••••••••••••••••	
	•••••		••••••
	••••••		•••••
(f)	The print indus	products of electrolysis of concentrated sodium chloride find extensive us stries. State the role of chlorine and sodium hydroxide in the paper industrials.	e in ry.
	(i)	Chlorine	(1 mark)
			••••••
			•••••
	(ii)	Sodium hydroxide	(1 mark)
			•••••
		•••••••••••••••••••••••••••••••••••••••	
		••••••••••••••••••••••••	

7.	(a)	(i)	Give the formulae of <b>two ionic</b> compounds that can be used to prepare sulphide salt.	lead(II) (1 mark)
			•••••••••••••••••••••••••••••••••••••••	••••••
			•••••••••••••••••••••••••••••••••••••••	
			***************************************	
		(ii)	Two moles of aqueous ammonia reacted with one mole of phosphoric(). Write an equation for the reaction that took place.	V) acid. (1 mark)
				• • • • • • • • • • • • • • • • • • • •
			•••••••••••••••••••••••••••••••••••••••	

(b) Solid copper(II) sulphate is available either as anhydrous or hydrated salt. Figure 4 shows enthalpy changes involved when water is added to each solid.




Figure 4

(i) Calculate the enthalpy change for the process:

	Water  CuSO <sub>4</sub>		(1 mark)

	(ii)	Describe how each of the following can be prepared starting w copper(II) sulphate.		ith aqueous
		I. hydrated copp	per(II) sulphate.	(2 marks)
		••••••••	······································	
		************		
		***************************************		
		II. anhydrous co	opper(II) sulphate.	(1 mark)
2				••••••
ř		***************************************		
		***************************************		
(c)	Alun	nium hydroxide is us	sed as an antacid.	
1067	(i) Name another compound that is used as an antacid.		(1 mark)	
		ranic another comp	Journa triat is used as an antacio.	
	(ii)	containing aluminiu	of hydrochloric acid in the stomach is 0.01 Normal hydroxide is used, calculate the mass of se 100.0 cm <sup>3</sup> of the stomach acid	M. If an antacid the antacid
		(A1 = 27.0; O = 16)	6.0; H = 1.0).	(3 marks)
		***************************************	······································	•••••
		***************************************		
		***************************************		••••••••••••••
		***************************************		***************************************
				***********
			······································	



#### THIS IS THE LAST PRINTED PAGE.