



SERIES 17 EXAMS

233/3

CHEMISTRY

PAPER 3

1. Procedure II

Table 1..... 3 marks

(a) – Complete table (C.T) ✓ 1 Mark

Conditions

- (i) Complete table with 3 titrations – 1 Mark
- (ii) Incomplete table with 2 titrations – ½ Mark
- (iii) Incomplete table with 1 titration – zero mark

Penalties:- Penalise ½ mark each

- (i) Wrong arithmetic
- (ii) Inverted table
- (iii) Burette reading beyond 50cm (unless explained)
- (iv) Unrealistic titre values i.e. 1cm³ or 100

(b) **Use of decimals** (tied to 1st and 2nd rows only)**Conditions**

(½ mark)

- (i) Accept 1 or 2 decimal points used consistently, if not penalise fully.
- (ii) Where 2 decimal points used the 2nd decimal point should be “0” or “5” if not penalise fully.
- (iii) Accept consistency in use of zero as initial burette reading i.e. 0, 0.0, 0.00

(c) **Accuracy** (Tied to correct titre value)..... (1 mk)

- (i) Atleast one of candidate’s values is within ± 0.1 of s.v (1mk)
- (ii) If non of candidates’ value is within ± 0.2 of s.v (0 mark)
- (iii) If one of the candidates value is within ± 0.2 of the s.v (½ mark)

(d) Principles of Averaging 1 mark

Conditions

- (i) - 3 consistent values averaged
- If 3 titrations done but only are consistent and averaged
- If 2 titrations done and are consistent and averaged

} 1 mark

Penalties

- Wrong arithmetic error is outside ± 0.2 units in d.p. ½ mark
- No working shown but answer is given correctly ½ mark
- Wrong workings with correct answer 0 mark

(e) Final accuracy (Tied to correct average titre)(1 mark)

Compare candidate’s average titre with the s.v

- i) If the candidates value is in ± 0.1of the s.v. – (1 mark)
- ii) If the candidate’s value is in ± 0.2 of the s.v. – (½ mark)
- iii) If the candidate’s value is beyond ± 0.2 – (0 mark)


beyond ± 0.2 0 mark

(ii) number of moles of the acid used

$$n = \frac{MV}{1000} = \left\{ 0.15x \frac{\text{Ans.in a (i)}}{1000} \right\} = \text{Ans } \underline{\hspace{2cm}} \text{ mole } \frac{1}{2} \text{ mark}$$

$$\text{e.g. } 0.15 \times \frac{20.35}{1000} = 0.003053 \text{ mole}$$

(b) Equation



States wrong/missing or letters joined ✓ ½ Mark

(c) (i) No. of moles of metallic carbonate in 25cm³ of Md.

$$= (\text{Answer in a (ii) above} \times \frac{1}{2})$$

= Answer _____ mole ✓ ½ Mark

e.g. $0.003053 \times \frac{1}{2} = 0.00153$ mole

(ii) No. of moles of the metallic carbonate in 50.00 cm^3 of solution Mc ✓ 1 Mark

50 cm^3 of Mc has same No. of moles of carbonate as 250 cm^3 of Md.

but 25 cm^3 of Md → Answer in c (i) above.

∴ 250 cm^3 of Md has Ans. c (i) $\times \frac{250 \text{ cm}^3}{25 \text{ cm}^3}$

= Ans. _____ mole ✓ ½ Mark

e.g. $\left(0.00153 \times \frac{250}{25}\right) = 0.0153$ mole

(iii) Molar mass of the metallic carbonate

50 cm^3 of Mc → Ans. c (ii)

80 cm^3 of Mc → ?

∴ Moles in 80 cm^3 of Mc = $\left(\text{Ans. c (ii)} \times \frac{80}{50}\right)$ moles

but 80 cm^3 of Mc has 7.0g

⇒ $\left\{\text{Ans c (ii)} \times \frac{80}{50}\right\}$ mole = 7.0g

1 mole - ?

So molar mass = $\left(\frac{1 \times 70}{\text{Ans c(ii)} \times \frac{80}{50}}\right)$

= Ans _____ g ½ mark

e.g. $= \frac{7.0 \times 1}{\left(0.0153 \times \frac{80}{50}\right)} = \frac{7.0}{0.02448} = 285.5477 \text{ g}$

(iv) Value of x in $\text{M}_2\text{CO}_3 \cdot x\text{H}_2\text{O}$

Let molar mass = $(23 \times 12) + 12 + (16 \times 3) + x(2 + 16)$

= $106 + 18x$

But molar mass = Ans. c (iii)

∴ $106 + 18x = \text{Ans. c (iii)}$

$x = \left\{\frac{\text{Ans.C (iii)} - 106}{18}\right\}$ ✓ ½ mark

= Ans _____ ½ mark

e.g. $106 + 18x = 285.5477$

$x = \frac{285.5477 - 106}{18}$

= $9.9749 \approx 10$

2. (a) Table ✓ 3 ½ marks

1. Complete table ✓ 1 Mark

Conditions

i. Complete table with 7 readings ✓ 1 Mark

Incomplete table with 5 – 6 readings ✓ ½ Mark

Incomplete table less than 5 readings 0 mark

ii. Treat initial value above 40°C and below 10°C as unrealistic and penalize ½ mark tied to $t = 0$

iii. Penalise ½ mark for each reading greater than 50°C from $t = 30$ seconds to a maximum of ½ mark.

iv. Penalize fully if all readings are constant.

2. Use of decimals ✓ 1 Mark

Accept whole numbers or readings with .0 or .5 used consistently, otherwise penalize fully.


3. Accuracy ½ mark

Compare the candidate's initial temperature (at time = 0) with the school value: If within ± 0.2 award 1 mark otherwise penalize fully

4. Trend ✓ 1 Mark

Award the first ½ mark for a continuous rise in temperature upto a maximum or constant values followed by a drop.

(b) Graph ✓ 3 Marks



Trend

Volume of R

(i) Labeling (both axis) ✓ ½ Mark

Penalize fully for – inverted axes

- wrong units

Accept if units are omitted

(ii) Scale ½ mark

Area covered by the plots should be at least ¾ of the plotting area: otherwise penalize fully.

(iii) Plotting ✓ 1 Mark

- Award ✓ 1 mark for at least 7 points correctly plotted

- Award ½ mark for 5 – 6 points correctly plotted otherwise award zero.

- Award fully for plots if the axes are inverted but the plotting is correct.

(iv) Shape 1 mark

- Award ½ mark for a straight line showing progressive increase in temperature.

- Award the other ½ mark for an extrapolated straight line showing a drop.

(c) (i) ½ mark - shown on the graph

(ii) ½ mark - value

(d) Heat change 1 ½ marks

$$\Delta H = Mc\Delta T$$

$$\text{e.g. } \frac{42.5}{1000} \times 4.2 \times 4.5 = -0.8033\text{Kj}$$

- Penalize ½ mark for wrong or absence of units

- Penalize ½ mark for the absence of the –ve sign on the answer

(e) Moles of NaOH = $\frac{25 \times 0.6}{1000} = 0.015\text{mol}$ ✓ ½ Mark

Molar enthalpy

$$0.015 \rightarrow -8033$$

$$1 \text{ mole} \rightarrow ?$$

$$= \frac{1}{0.015} \times .8033 = -53.5533 \text{ kJ mol}^{-1} \quad \checkmark \quad \frac{1}{2} \text{ Mark}$$

2. (II)

1. (a) Complete table ✓ 5 Marks

(i) Complete table with 10 readings 2 ½ marks

- Penalize ½ mark for each space not filled

- Penalize ½ mark for any time reading less than or greater than 220 seconds.

- If the candidate enters some readings in fractions and others in decimals award accordingly (each value is worth ½ mark)

(ii) Use of decimals ½ mark

Accept whole numbers or 1dpl – 2dpl used consistently throughout: otherwise penalize fully.


(iii) Accuracy ½ mark

Compare the candidate's first reading and the school value. If within ± 2 seconds award ½ mark: otherwise penalize fully.

(iv) Trend ½ mark

Award ½ mark if time increasing throughout: otherwise penalize fully.

(b) The graph



(i) Scale ½ mark

Area covered to be ¾ of the space provided. Scale should accommodate 4 plots, otherwise penalize fully.

I	Observations	Inferences
	Slow effervescence ✓ 1 mark/ fizzing formation of bubbles of colourless gas Accept: Slow production of colourless gas Rej: "hissing" ½ mark	- $H^+/H_3O^+/-COOH/R - OH$ ✓ 1 mark - Carboxylic group in words - Solution is acidic Rej: Solution is an acid ½ mark
II	- Orange colour of $H^+/K_2Cr_2O_7$ persists/is retained Rej: Colour of $H^+/K_2Cr_2O_7$ retained ½ mark	Absence of $R - OH$ ½ mark
III	Red/orange/yellow bromine water decolourised ✓ 1 mark Rej: Bromine water turns colourless /dicoloured ✓ ½ mark	$\begin{matrix} \diagdown & & \diagup \\ & C = C & \\ \diagup & & \diagdown \end{matrix}$ or $-C \equiv C -$ ✓ 1 mark Accept: Unsaturated organic compound Penalise fully for any contradictory functional group ✓ ½ mark
IV	Method - Place $1cm^3$ of solution T in a test tube - Add 1 – 3 drops of universal indicator solution - Match the colour obtained with the pH chart pH value 4 Rej: pH range of 4 – 5 ✓ ½ mark	Solution is weakly acidic Rej: Weak acid ✓ ½ mark