MARKING SCHEME

CHEMISTRY PRACTICAL

	I	II	III
Final burette reading (cm³)			
Initial burette reading (cm³)			
Volume of solution T used (cm³)			

a) Complete table ………………………………………………………………….(1mk)
Complete table with 3 titrations done – 1mk
In Complete table with 2 titrations done - 1mk
incomplete table with 1 titration done – 0mks

Penalize ½ mk once for
➢ Inverted table
➢ Wrong arithmetic
➢ Unrealistic titre values (below 1 or above 50 unless explained)

b) Use of decimals……………………………………………………………………..1mk
➢ Accept 1 or 2 d.p uses consistently otherwise penalize fully
➢ If 2 d.p used the 2^{nd} d.p should be either 0 or 5 otherwise penalize fully
➢

c) Accuracy …………………………………………………………………….1mk
Compare the candidates titre values with the S.V
➢ If any value is within +- 0.1 award   1mk
➢ If within + -0.2 award ½ mk
➢ If beyond +- 0.2 award zero mark

d) Principles of averaging ……………………………………………………….1mk
1f 3 consistent titrations done and averaged   1mk
If 3 titrations done but only 2 are consistent and averaged   (1mk)

If only two titrations done, are consistent and averaged   (1mk)
If 3 titrations done and are consistent but only 2 are averaged   (0mk)
If 3 inconsistent titres averaged   (0mk)
1f 2 inconsistent titres averaged   (0mk)

e) Final answer accuracy…………………………………………………………..1mk
Compare the candidates correct average titre with S.V
➢ If within +- 0.1 of S.V   (1mk)
➢ If within +- 0.2 of S.V   (½ mk)
➢ If beyond +- 0.2 of S.V   (0mk)

a) Calculate the:

    i.   Average volume of solution T used   (1 mark)

Captured in principal of averaging

    ii.   The number of moles of solution Z used in titration   (1 mark)

$$\frac{25.0 \times 0.4}{1000} = 0.01 \ moles$$

    iii.   Concentration of solution T in moles per liter.   (2 marks)

$$NaOH_{(aq)} + HCl_{(aq)} \longrightarrow NaCl_{(aq)} + H_2O_{(l)}$$

0.01 moles $\longrightarrow$ Average titre Volume

$\longleftarrow$ 1000cm³

$$\frac{1000 \times 0.01}{Average \ titre \ Volume}$$

    iv.   Calculate the mass of sodium carbonate that reacted with solution Y. (Na= 23, C= 12, O= 16)   (3marks)

Calculate the mass of sodium carbonate that reacted with solution Y. (Na= 23, C= 12, O= 16)   (3marks)

$$Na_2CO_{3(s)} + 2HCl_{(aq)} \longrightarrow NaCl_{(aq)} + CO_{2(g)} + H_2O_{(l)}$$

Number of moles of HCl in average titre $\longrightarrow$ 0.01moles

250cm³ $\longrightarrow$?

$$\frac{250 \times 0.01}{Average \ titre \ Volume}$$

Original number of moles of HCl = $\frac{2 \times 80}{1000}$ = 0.16moles

Moles of HCl that reacted with Carbonate = 0.16 − number of mol in 250cm³

Moles of Sodium Carbonate = ½ × the difference in moles aba...

RFM of Na₂CO₃ = (23×2) + 12 + (16×3) = 106

106g $\longrightarrow$ 1mole

? $\longleftarrow$ No. of moles of sodium Carbonate

Mass of Sodium Carbonate = Number of moles of Na₂CO₃ × 106

                                     (1 mark)

v.   Work out the percentage purity of solid X.   (1 mark)

$$\frac{mass\ of\ sodium\ carbonate\ that\ reacted\ with\ solution\ Y}{5} \times 100$$

b) Write the ionic equation between solid X and solution Y.   (1 mark

$$2H^+_{(aq)} + CO_3^{2-}_{(aq)} \longrightarrow CO_{2(g)} + H_2O_{(l)}$$

3. **You are provided with solid L. Carry out the test below and record your observations and inferences in the spaces provided.**

a) Describe solid.   (2 marks)

b) Place half spatula of L in a non-luminous flame of a Bunsen burner.

Observations	Inference
Solid melts and burns with a yellow sooty flame ($\frac{1}{2}$ mark)	$C=C$, $-C\equiv C-$ ($\frac{1}{2}$ mark)

c) Dissolve the remaining solid L in the distilled water and divide the solution into four portions.

i.   To the fourth portion, add few drops of acidified potassium chromate (vi) and warm.

Observations	Inference
Orange acidified potassium chromate VI does not change to green *Reject persist or remains or no color change, no observable change* ($\frac{1}{2}$ mark)	R-OH absent ($\frac{1}{2}$ mark)

ii.   To the second portion, add few drops of bromine water and warm.

Observations	Inference
Yellow bromine water changes to colorless / decolorized (**¹/₂ mark**)	$\overset{\mid}{C}=\overset{\mid}{C}$, $-C\equiv C-$ (¹/₂ mark)

iii.   To the third portion add few drops of acidified potassium manganite (vii) and warm.

Observations	Inference
Purple acidified potassium manganate VII changes to colourless (¹/₂ mark)	$\overset{\mid}{C}=\overset{\mid}{C}$, $-C\equiv C-$ (¹/₂ mark)

iv.   To the fourth portion add a quarter spatula end-full of sodium hydrogen carbonate.

Observations	Inference
Effervescence / bubbles produced (¹/₂ mark)	R-COOH (¹/₂ mark)