

**233/1**

**CHEMISTRY**

**PAPER 2**

**THEORY**

**TIME – 2HRS**

**SUKELLEMO PRE MOCK JOINT EXAMS**


Pre Mock Examination

JUNE 2022

**MARKING SCHEMES**

- a) i) To remove dust particles ✓ - than would otherwise “poison” that catalyst (1mk)  
 ii) 9 atmospheres ✓ (1mk)  
 iii) To pre heat NH<sub>3</sub> and air to an optimum temperature (reactants) ✓  
 To cool NO (product) ✓ (2mks)  
 iv) Platinum –Rhodium catalyst ✓ (1mk)  
 v) I :  $4\text{NH}_3(\text{g}) + 5\text{O}_2(\text{g}) \longrightarrow 4\text{NO}(\text{g}) + 6\text{H}_2\text{O}(\text{g})$  ✓ (1mk)  
 II :  $2\text{NO}(\text{g}) + \text{O}_2(\text{g}) \longrightarrow 2\text{NO}_2(\text{g})$  ✓ (1mk)  
 III  $4\text{NO}_2(\text{g}) + \text{O}_2(\text{g}) + 2\text{H}_2\text{O}(\text{l}) \longrightarrow 4\text{HNO}_3(\text{aq})$

b) i)



(3mks)

Moles of O<sub>2</sub> ⇒  $\frac{1}{2} \times 0.25$   
 = 0.125 ✓  $\frac{1}{2}$


Volume of O<sub>2</sub> ⇒  $0.125 \times 22.4$   
 = 2.8 dm<sup>3</sup>

V = 280 cm<sup>3</sup>

- ii) Manufacture of fertilizers ✓  
 Manufacture of explosives ✓  
 Manufacture dyes and drugs ✓ (2mks)  
 Purification of metals eg. silver and gold  
 Etching designs on some metals

2. (a) this is the maximum mass of salt/solute that can saturate 100g of water at a given temperature.

(b)



(c) (i) 30g of H<sub>2</sub>O     Accept 36g/100g water

(ii) 33.5°C

(d) (i) Sol at 40°C = 61g/100g H<sub>2</sub>O

At 27°C

(ii) Sol at 40°C = 61g/100g H<sub>2</sub>O

Sol at 27°C = 35g/100g H<sub>2</sub>O

Sol at 5°C = 10g/100g of H<sub>2</sub>O

(e) Y is more soluble than Z.

Y dissolves better at higher temperatures than Z

(f) Fractional crystallization – Separation of mixtures with different solubilities

3. (a) (i) I:  $Zn(OH)_2$  ✓<sup>1</sup>  
 II:  $ZnCl_2$  ✓<sup>1</sup>  
 III:  $ZnO$  ✓<sup>1</sup>
- (ii)  $Pb_{(aq)}^{2+} + 2Cl_{(aq)}^- \rightarrow PbCl_{2(s)}$  ✓<sup>1</sup>
- (iii) White precipitate soluble in excess. ✓<sup>1</sup>
- (b) Ammonia gas is polar and ionizes ✓<sup>1</sup> in water which is polar.  
 While it does not ionize in methylbenzene which is non polar.
- (c) (i) Calcium carbonate/magnesium carbonate.  
 (ii) Passing a solution of dilute hydrochloric acid or nitric (V) acid  
 in the boiler. (1mk)
- (d)  $[Zn(OH)_4]^{2-}$

4a

- i) Hygroscopy
- ii) Deliquescence
- iii) Efflorescence



| Fe   | S    | O    | H <sub>2</sub> O             |
|------|------|------|------------------------------|
| 20.2 | 11.5 | 23.0 | 45.3                         |
| 56   | 32   | 16   | 18                           |
| 0.36 | 0.36 | 1.44 | 2.52 $\checkmark 1\text{mk}$ |
| 1    | 1    | 4    | 7                            |

$$(\text{FeSO}_4 \cdot 7\text{H}_2\text{O}) = 278$$

$$278n = 278$$

$$n = 1 \checkmark 1\text{mk}$$

$$\text{Formula FeSO}_4 \cdot 7\text{H}_2\text{O} \checkmark 1\text{mk}$$

ii) No. of moles  $\frac{6.95}{278} = 0.025 \text{ moles} \checkmark 1\text{mk}$

$$0.025 \text{ moles} \cdot 250 \text{ cm}^3$$


$$1000$$

$$\frac{0.025 \times 1000}{250} = \frac{25}{250} = 0.1 \text{ mole/litre} \checkmark 1\text{mk}$$

d)

- Add excess lead carbonate to dilute HNO<sub>3</sub>,  $\checkmark 1\text{mk}$
- shake and filter to remove unreacted carbonate  $\checkmark 1/2\text{mk}$
- Add excess dilute HCl to the mixture  $\checkmark 1/2\text{mk}$
- Filter to obtain lead(II) chloride as the residue  $\checkmark 1/2\text{mk}$
- Rinse and Dry between filter paper to obtain solid PbCl<sub>2</sub>  $\checkmark 1/2\text{mk}$

5. i)



ii) – Catalyst Nickel  
 - Temperature 150°C to 250°C

iii) magnesium is very expensive  
 it gives a mixture of gases including bad smelling and poisonous hydrogen

sulphide

$\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$

v) – Manufacture of ammonia  
 - Manufacture of hydrochloric acid  
 - Welding and cutting of metals  
 - Rocket fuel and in fuel cells.

b) i) Slowly to allow ample time for reaction repeatedly to ensure all active air (oxygen) is used up  
 ii) The brown copper turnings slowly changed black.

iii)

$$\frac{120 - 95.5}{120} \times 100$$

$$\frac{24.5}{120} \times 100 = 20.41667\%$$


6. (a) B – 2 : 8  $\sqrt{\frac{1}{2}}$

D – 2 : 8 : 8  $\sqrt{\frac{1}{2}}$


(b) (i) D  $\sqrt{\frac{1}{2}}$

(ii) E  $\sqrt{1/2}$

(c) Atomic radius of B is larger  $\sqrt{1}$  than that of C. C has more protons. The outer energy level electrons are pulled  $\sqrt{1}$  strongly to the nucleus reducing the atomic size.



(e) Element B has stronger metallic  $\sqrt{1}$  bond ( has more delocalized electrons ) than A, hence higher amount of heat  $\sqrt{1}$  energy is needed to break the bond.



- Reject fully if unbalanced

- Award  $\frac{1}{2}$  mk if states are missing or any one state is wrong.

(g) Add water  $\sqrt{1/2}$  to the mixture and stir.

Filter  $\sqrt{1/2}$  to obtain lead (II) sulphate as  $\sqrt{1/2}$  residue and sulphate of E as filtrate

Dry the residue  $\sqrt{1/2}$  to obtain lead (II) sulphate.

Evaporate  $\sqrt{1/2}$  the filtrate to dryness  $\sqrt{1/2}$  to obtain the solid sulphate of E.

a) Ethylpropanoate

b) But-2,3-diene

- b) i) A - yeast ✓ (1mk)  
 I - fractional distillation (1mk)  
 B - Sodium ethoxide (1mk)  
 C - Hydrogen (1mk)  
 D - Ethene (1mk)  
 E - polyethene / polythene (1mk)

(4mks)

iii) I :  $C_6H_{12}O_6 \rightarrow 2C_2H_5O + 2CO_2$  ✓ (1mk)  
 $mole\ of\ sugar \frac{144000}{180} = 800\ moles$

$$mole\ of\ ethanol = 2 \times 800$$

$$= 1600\ moles$$

II (3mks)

$$nfm \rightarrow 2 \times 12 + 6 + 16 = 46$$

$$mass\ of\ ethanol = \frac{46 \times 1600}{1000}$$

$$= 73.6\ kg$$